Identical Particles

Indiscernibility of Identicals

$$\forall x = y \rightarrow \forall P (Px \leftrightarrow Py)$$

2nd order modal logic

Identity of Indiscernibles

$$\forall P (Px \leftrightarrow Py) \rightarrow x = y.$$

(Leibniz principle)

weak ~~strong~~ version 'P' includes location
~~weak~~ strong 'P' as it includes location.

In classical mechanics weak version at least
(but not strong version due to impenetrability)
is violated we can have 2 distinct objects.
they are indistinguishable and yet
numerically distinct (i.e. not identical)

So Identity ≠ Indiscernibility
But Indiscernibility ≠ Identity.

This is to treat classical particles
particles are individuals if they are
distant 'things' persisting identical
individuals — they can be described
under names e.g. John ≠ John.

Distinct But can use language predicates
at a time — John at t1 unique object having
the parts that John has
But he can at a time spatially disjoint

event of making the choice possible.

Irreducible uncertainty is tied up in the Copenhagen attitude to doing stat. mechanics

a)

different callum/row b)

4 possible states and equal weights

c) 

d)

So classes d, f are probs and each $\frac{1}{2}$.

But in QM the rule for Born probs is 1/3

here. a) & d) are counted in the same rule.

Does this mean that in QM. electrons do not possess theoretical individuality?

- they are in same pure individuals.

It is often claimed that this is that electrons are non-individuals because of the Pauli principle as.

we shall see that distinction in terms of individuals & non-individuals is available both in classical and Q. mechanics

an entity that (ontologically)

Note An individual can be identified, as this object rather than that.

A. Continuant can be (ontologically) identified as the same individual in virtue of its

Two objects are identical if they are (ontologically) the same object.

~~A particle~~ An individuator is that feature of an individual which individuates it

Two objects are indistinguishable if they have all the same properties.

A particle is a continuant (which is individuated by its individuality) (transcendent solipsistic its properties) and which has a non-zero location which varies over time, but no extension (point particle)

A field configuration is an individual which is a non-continuant (0- or 1-dimensional) which is individuated by the transcendent individuality of its spatial locations.

When 2 ~~particles~~ indistinguishable particles change places there is ontologically a new state of affairs.

When 2 indistinguishable flips in a field change place.

no new state is created for

Individual A → Individual B

" B → " A.

In a fluid, a velocity (as this is a particle here) is being passed from one molecule to another (i.e. from one place to another)

10. In non-contiguous an individual can change into another included at various times.

This is what happens when a flea passes on a field.

Included A becomes included B as particle passes.

The large included moves from one location to another in one day as instead probably for another.

N.B. There cannot be more than one included object.

Not There can be 2 or more individuals which are contiguous as the 2 or more individuals which are non-contiguous.

This applies the question of how to can have a they of Nat. Included (only non-contiguous) and needs to be how many than one of them.

But note that ephemerals may also be regarded as

relative continuants which can be reidentified through time as the same S, for some total predicate S.

In the case of a fluid flow and can identify a as the same flow as it was stated to persist continuity in the development of flow to flow a

If you cannot step in the same river twice (Heraclitus).
You can step in the same river, but not the same water

A river is a relatively persistent a flow of water as an (abstract) continuant.

Note that while a fluid configuration is as observed, a fluid system may be regarded as the changing state of a set of continuants, the primary individuals of the flow - time 'fluids'

A process is a succession of states of affairs involving different continuants but showing some invariant features S a is the same S as it is a relatively identity statement, A process will be need to persist

Persistents { continuants — resistance of matter
non-identifiable ephemerals — resistance of form

→ micro-things

ex plane, cell, River

Theseus' ship

process-things exhibit relative identity

'a' is same as 'b'

partial predicate S.S.

1986 Seminars Individuals are physical particulars

Continuants (particles and places?) → Hyperon decays

ephemerals (process stages instants of time)

- persistent ephemerals (process-things) eg. planes resistance of resistance of form
- non-persistent i.e. non-identifiable ephemerals. like mixed-up waves.

Entities which are individuals but not continuants I shall call *eponerals*

Note that flaps in a fold are eponerals even if they do not merge if no record. Movement of an eponeral or its detachment at one point and creation at another point.

Ex. Even if the two parts do coincident this mode of description makes sense.

The change of state of an eponeral is described by an equation $\delta = \xi\eta$. where η destroys the eponeral in its original location and ξ records it in its new location.

If a continuant δ only changes its location in one (fixed under) it persists in $\delta\xi$ state too.

The change in description of change. from a description in terms of continuants to a description in terms of eponerals corresponds to the change from δ to the factorizing form $\xi\eta$.

Here is just a tool in classical mechanics as in quantum theory in the transition from 1st quantization to 2nd quantization



the movement of a particle from A to B
is decided by the poles.

$$\xi(A) = B = \xi_B \eta_A$$



We can describe elementary particles either as continuants or as ephemerals.

Let us opt where particles do get created and destroyed. The description in terms of ephemerals becomes much more apposite.

yes/no (declarative field) for questions of an ephemeral in terms of occupation of space. We can introduce labels (value names) neg. continuants of space-time regions in the field description. The zeros can we can get exist counter to classical Stapledon / nucleus stable in



cf Resources of Nature regions of configuration space. See Weekes in 'Models, Foundations of Physics' p. 139 ff. in Hall, Hall 'Unpublished Papers of Isaac Newton'.

N-partide S Eq. ⟶ Field quantized. → 'field's

1-particle S. Eq. ⟶ 2nd quantization → quantized field.

Field system ⟶ Field quantization → quantized field.